



VALVULAR HEART DISEASE

LONG TERM DURABILITY OF EDWARDS BALLOON EXPANDABLE TRANSCATHETER AORTIC VALVE IMPLANTATION

ACC Poster Contributions

Georgia World Congress Center, Hall B5

Monday, March 15, 2010, 3:30 p.m.-4:30 p.m.

Session Title: Outcome analysis in Valvular Disease

Abstract Category: Valvular Disease

Presentation Number: 1227-390

Authors: Ronen Gurvitch, Edgar Tay, Anson Cheung, Fabian Nietlispach, Jian Ye, David A. Wood, Chris Thompson, Namal Wijesinghe, Jonathon Leipsic, Samuel V. Lichtenstein, John G. Webb, St. Paul's Hospital, Vancouver, BC, Canada

Background: Whilst Transcatheter Aortic Valve Implantation (TAVI) using the Edwards Balloon Expandable Valve (EBEV) has been shown to be safe and effective, no data is available on long-term valvular function and clinical outcomes.

Methods: To assess long-term durability, we analysed all patients who underwent successful TAVI who survived >30 days and reached a minimum of 3 years follow-up. Clinical follow-up, echocardiographic and Computed Tomography (CT) scanning were performed to assess valvular function.

Results: 69 patients were analysed at a median of 3.4 years (IQR 3.2-3.8). Longest follow-up was 4.7 years. Baseline risk was high (mean logistic EuroSCORE 31 ± 15 , Society Thoracic Surgery score 9.0 ± 3.5), mean age 83.4 ± 7.5 years. Survival at a median of 3.4 years was 64%. The most common mortality cause was respiratory (30%). There was no mortality related to valvular dysfunction or failure. There were no cases of valve failure or repeat operation/intervention. Functional class improvements, Aortic Valve Area (AVA) and mean transaortic gradients were maintained long-term (see table). In the 10 patients who survived >4 years to date, the mean increase in transaortic mean pressure from 12 to 48 months was 3.0 ± 2.1 mmHG ($p < 0.05$), and mean AVA decrease of 0.23 ± 0.16 cm² ($p < 0.05$). In 11 patients who underwent CT follow-up, no evidence of structural deterioration, fracture or recoil were detected.

	Baseline	1 year	2 years	3 years
NYHA Class †	3.0 ± 0.56	1.3 ± 0.56	1.3 ± 0.48	1.4 ± 0.56
AVA, (cm ²) †	0.59 ± 0.16	1.50 ± 0.34	1.46 ± 0.36	1.42 ± 0.27
AVMG, mmHG †	45.2 ± 4.7	11.4 ± 5.6	12.8 ± 4.9	12.6 ± 5.0
NYHA=New York Heart Association functional class, AVA = Aortic Valve Area, AVMG=Aortic Valve Mean Gradient, † mean \pm SD				

Conclusions: TAVI with the EBEV shows durability to over 4 years with no evidence of structural failure.